



**HAL**  
open science

## **Are trait self-control and self-control resources mediators of relations between executive functions and health behaviors?**

Cyril Forestier, Margaux de Chanaleilles, Roxane Bartoletti, Boris Cheval, Aïna Chalabaev, Thibault Deschamps

### ► **To cite this version:**

Cyril Forestier, Margaux de Chanaleilles, Roxane Bartoletti, Boris Cheval, Aïna Chalabaev, et al.. Are trait self-control and self-control resources mediators of relations between executive functions and health behaviors?. *Psychology of Sport and Exercise*, 2023, pp.102410. 10.1016/j.psychsport.2023.102410 . hal-04061618

**HAL Id: hal-04061618**

**<https://nantes-universite.hal.science/hal-04061618v1>**

Submitted on 11 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Journal Pre-proof

Are trait self-control and self-control resources mediators of relations between executive functions and health behaviors?

Cyril Forestier, Margaux de Chanaleilles, Roxane Bartoletti, Boris Cheval, Aïna Chalabaev, Thibault Deschamps



PII: S1469-0292(23)00034-1

DOI: <https://doi.org/10.1016/j.psychsport.2023.102410>

Reference: PSYSPO 102410

To appear in: *Psychology of Sport & Exercise*

Received Date: 14 July 2022

Revised Date: 17 February 2023

Accepted Date: 3 March 2023

Please cite this article as: Forestier, C., de Chanaleilles, M., Bartoletti, R., Cheval, B., Chalabaev, Aï., Deschamps, T., Are trait self-control and self-control resources mediators of relations between executive functions and health behaviors?, *Psychology of Sport & Exercise* (2023), doi: <https://doi.org/10.1016/j.psychsport.2023.102410>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd.

Are trait self-control and self-control resources mediators of relations between executive functions and health behaviors?

Cyril Forestier<sup>1\*†</sup>, Margaux de Chanaleilles<sup>2†</sup>, Roxane Bartoletti<sup>3</sup>, Boris Cheval<sup>4,5</sup>,  
Aïna Chalabaev<sup>2</sup>, Thibault Deschamps<sup>1</sup>

<sup>1</sup>Laboratoire Motricité, Interactions, Performance, MIP - EA4334, Le Mans Université, Nantes Université, Le Mans, Nantes, France

<sup>2</sup>Univ. Grenoble Alpes, SENS, Grenoble, France

<sup>3</sup>LAPCOS, Université Côte d'Azur, Nice, France

<sup>4</sup>Swiss Center for Affective Sciences, University of Geneva, Geneva, Switzerland

<sup>5</sup>Laboratory for the Study of Emotion Elicitation and Expression (E3Lab), Department of Psychology, University of Geneva, Geneva, Switzerland

\*Corresponding author: 23, rue du Recteur Schmitt Bât F0 - BP 92235 44322 NANTES Cedex 3, France; cyril.forestier@univ-nantes.fr; @CForestier\_PhD (C. Forestier)

†C.F. and M.d.C contributed equally and share first authorship

Word count Abstract: 221

Word count text: 5843

Tables: 1

Figures: 3

### **Funding**

M.d.C. is supported by a PhD grant from Univ. Grenoble Alpes.

R.B. is supported by a Ph.D grant from Univ. Côte d'Azur.

B.C. is supported by an Ambizione grant (PZ00P1\_180040) from the Swiss National Science Foundation (SNSF).

### **Competing interests**

The authors declare no conflict of interests.

### **Consent for publication**

All the authors listed in the by-line have agreed to the by-line order and to the submission of the manuscript in this form.

### **Contributors**

C.F., M.d.C and T.D. designed the study protocol. C.F. designed the analyses. M.D. C.F, M.d.C, R.B, and T.D. collected the data. C.F. analyzed the data. C.F., M.d.C and T.D. drafted the manuscript. A.C., R.B, M.B, and B.C. critically appraised the manuscript and significantly contributed to its content and structure. All authors approved the final version of the manuscript.

### **Data sharing**

Data and code for the analysis are available on OSF at

[https://osf.io/hpsjw/?view\\_only=6e28c8307294494e9eec45d2670efd8d](https://osf.io/hpsjw/?view_only=6e28c8307294494e9eec45d2670efd8d)

1

**Abstract**

2 This study investigated associations between executive functions (i.e., inhibition, working  
3 memory, cognitive flexibility) and individual differences in self-control and health behaviors.  
4 We examined whether executive functions predict physical activity, sedentary activity, and  
5 healthy and unhealthy diets, and whether trait self-control and self-control resources mediate  
6 these associations. Three hundred and eighty-five participants completed a questionnaire  
7 assessing trait self-control and self-control resources, physical activity, sedentary activity, and  
8 healthy and unhealthy diets. They also performed three randomly ordered cognitive tasks, a  
9 stop-signal task (i.e., inhibition), a letter memory task (i.e., updating), and a number-letter  
10 task (i.e., switching). Structural equation modeling revealed that self-control resources  
11 predicted positively physical activity ( $R^2 = .08$ ), negatively sedentary activity ( $R^2 = .03$ ) and  
12 positively healthy diet ( $R^2 = .10$ ). Moreover, trait self-control predicted positively healthy diet  
13 ( $R^2 = .10$ ) and negatively unhealthy diet ( $R^2 = .19$ ). Moreover, analyses revealed that  
14 switching significantly predicted self-control resources, and highlighted three totally mediated  
15 relations between this executive function and physical activity, sedentary activity and healthy  
16 diet. However, no evidence was found supporting associations between inhibition and  
17 updating, and health behaviors, or relations mediated by self-control for these executive  
18 functions. The findings suggest the importance of trait self-control and self-control resources  
19 for health behavior adoption and pave the way for studies exploring the role of the executive  
20 functions in an affective context. Open materials [<https://osf.io/hpsjw/>].

21

22 Keywords: self-control resources, trait self-control, inhibition, updating, switching, health  
23 behaviors

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

## 1. Introduction

According to the World Health Organization, physical inactivity and an unhealthy diet are among the most important risk factors for noncommunicable diseases, causing one death every seven seconds and one death every three seconds, respectively (Forouzanfar, Afshin, Alexander, Biryukov, et al., 2016; Lee et al., 2012). Changing these unhealthy behaviors by improving the regularity of physical activity and healthiness of diet could prevent 16 million premature deaths each year (Forouzanfar, Afshin, Alexander, Anderson, et al., 2016). Despite widespread declarations of intention to adopt healthy behaviors, most people fail to reach minimum recommendations (Ford et al., 2011). In this context, some promising conflict resolution models could effectively promote health behaviors (Sniehotta et al., 2014).

Conflict resolution models (e.g., the integrative self-control theory, Kotabe et Hofmann, 2015; the goal-conflict model, Stroebe, 2022) emphasize that health-behavior facilitators (e.g., health-behavior goals, such as an intention to do more physical activity) face barriers (e.g., temptation toward the competing behavior, such as a desire to engage in a sedentary activity), leading to motivational conflicts that need to be resolved (Forestier, et al., 2022a; Rabiau et al., 2006). An adaptative resolution of this conflict (i.e., one in favor of the goal) promotes the health-behavior goal (e.g., going for a run, snacking on an apple). A maladaptive one promotes the competing behavior (e.g., remaining on the couch, snacking on a chocolate bar) (e.g., Gillebaart et al., 2016). Here, self-control is defined as the self-regulation operationalization, by which an individual resolves a motivational conflict, through either an effortless or effortful strategy (Forestier et al., 2022b; Gillebaart, 2018). As such, it is expected to play a key role in promoting the resolution of motivational conflicts, favoring healthy behaviors. Indeed, some studies showed correlations between self-control and healthy diet and weight control with a small-to-medium effect size (de Ridder et al., 2012,  $r = .17$ ), and self-control and physical activity (Pfeffer & Strobach, 2018,  $r = .29$ ).

26 In 2018, Forestier et al. identified individual differences in trait self-control and self-  
27 control resources, as two dimensions independently associated with health behaviors. Despite  
28 their importance, self-control resources have been mostly considered in ego-depletion  
29 research (e.g., Rouse et al., 2013; see Forestier et al., 2022 for more details) and rarely on  
30 research on self-control and health behaviors (de Ridder et al., 2018). However, defined as  
31 “the objective and subjective amounts of energy available for the self to initiate a self-control  
32 act” (Forestier et al., 2022b, p. 21), the self-control resources seem crucial for motivational  
33 conflict resolution, and can be assessed through subjective perception of energy availability,  
34 or physiological markers (e.g., vagal activity). Accordingly, trait self-control distinguishes  
35 individuals with a more or less tendency to successfully resolve motivational conflicts. In  
36 parallel, individual differences in self-control resources distinguish individuals with a more or  
37 less tendency to experience a high level of self-control resources (Forestier et al., 2018).  
38 Authors identified that a remarkable portion of variance (40%) in self-control resources is  
39 found at the between-person level (Smolders et al., 2013). In short, individuals with high trait  
40 self-control consume a healthier diet than those with low trait self-control; individuals with  
41 high self-control resources practice more physical activity, are less sedentary, and eat  
42 healthier than individuals with low self-control resources. These results are in line with Finne  
43 et al. (Finne et al., 2019), highlighting that within-person self-control fluctuations are more  
44 important than between-person differences to predict physical activity. Moreover, a recent  
45 study showed that within-person variations of self-control strength (i.e., a scale that could  
46 refer to what Forestier et al., (2022) proposed as “self-control willingness”) predict physical  
47 activity but not trait self-control (Englert & Rummel, 2016).

48 Despite these interesting results, Forestier et al. (2018) did not specifically examined  
49 the predictors of individual differences in the two self-control dimensions. Some hypotheses  
50 on the executive functions have been already put forward that can partly explain the

51 differences in these dimensions (e.g., Hofmann et al., 2012, Table 2). Specifically, inhibition,  
52 updating and switching (also called “inhibitory control”, “working memory”, and “cognitive  
53 flexibility” respectively) is likely related to self-control, and individuals with higher executive  
54 abilities would, therefore, be good self-controllers (Friese et al., 2011; Hofmann et al., 2011).  
55 Moreover, executive functions could operationalize the “self-control capacity” that makes  
56 self-control success possible (Forestier, et al., 2022b; Kotabe & Hofmann, 2015). Presumably,  
57 for instance, individuals with high inhibition capacity should present high self-control through  
58 a better ability to override prepotent responses such as habits or impulses that are  
59 incompatible with goals attainment. Likewise, individuals with high working memory  
60 abilities could better maintain information in an active, quickly retrievable state and shield  
61 this information from distraction. They are therefore better able to use goal-relevant  
62 information. Finally, relative to individuals with lower switching ability, individuals with high  
63 switching may present high self-control through a high flexibility (opposed to “rigidity”),  
64 which allow goal attainment by abandoning suboptimal means and selecting alternative  
65 means to reach the same goal (see Hofmann et al., 2011, 2012, for full theoretical discussion).  
66 Additionally, structurally and functionally, brain networks in the prefrontal cortex that control  
67 executive function, overlap with networks involved in self-control (e.g., right inferior frontal  
68 gyrus, Cipolotti et al., 2016; Lopez et al., 2016).

69         Several assumptions on how executive functions and self-control might interact have  
70 been proposed, including the mediation model hypothesis (Hofmann et al., 2012). To date,  
71 most empirical self-control studies examined executive functions as moderators of the relation  
72 between self-control and health behaviors (Hofmann et al., 2009a; Pfeffer & Strobach, 2017),  
73 with mitigated results. Hence, we currently aimed at addressing the rarely explored approach  
74 of executive functions as direct predictor of self-control. As far as we know, only Saunders et  
75 al. (2018) and Necka et al. (2018) have investigated the association between executive



76 functions and self-control. Saunders et al. (2018) found no evidence of a correlation between  
77 trait self-control and inhibition. Using structural equation modeling, Necka et al. (2018) found  
78 no significant association between a “trait self-control” latent variable and an “executive  
79 functions” latent variable. However, methodological improvement would allow some results-  
80 related pitfalls to be avoided. First, only the relations between latent variables were tested,  
81 without examining the independent contribution of each executive function to trait self-  
82 control. Second, these previous studies never focused on self-control resources, which are  
83 correlated with trait self-control but remain an independent dimension to be considered  
84 (Forestier et al., 2018). Third, the possibility that trait self-control and executive functions are  
85 associated with health behaviors was not investigated. Yet, the relation between executive  
86 functions and health behaviors deserves attention. Indeed, another study showed that  
87 individuals with the highest inhibition adopted a less unhealthy diet (Hofmann et al., 2009).  
88 Similarly, high updating abilities have been associated with more physical activity  
89 (Lambourne, 2006; Pfeffer & Strobach, 2017). Finally, a bi-directional relationship between  
90 executive functions and health behavior has been proposed, with individuals with high  
91 executive abilities being more likely to adopt a healthy lifestyle that would, in turn, enhance  
92 their executive functions in the long run (Allan et al., 2016). Empirical data has recently been  
93 reported supporting this relation (Cheval et al., 2020). Accordingly, if executive functions are  
94 correlated with self-control, as advanced theoretically (Hofmann et al., 2012), they could  
95 promote health behaviors through direct and indirect effects, partially mediated by trait self-  
96 control and self-control resources. Such assumptions have never yet been tested.

97         The current study aimed to investigate the independent contribution of each executive  
98 function to self-control and four crucial health behaviors to prevent noncommunicable  
99 diseases, while considering differences among individuals in their trait self-control and self-  
100 control resources. For these purposes, structural equation models were used, one for physical

101 activity versus sedentary activity, and a second for a healthy versus an unhealthy diet, as in  
102 Forestier et al., (2018). In line with theoretical discussions (Hofmann et al., 2012), we  
103 hypothesized that executive functions would positively predict trait self-control and self-  
104 control resources (H1) and healthy behaviors (physical activity and healthy diet) (H2), and  
105 negatively predict unhealthy behaviors (sedentary activity and unhealthy diet) (H3). Similarly,  
106 we hypothesized that trait and self-control resources would positively predict healthy  
107 behaviors (H4) and negatively predict unhealthy behaviors (H5). Finally, the mediated  
108 relation implies that (H6) executive functions will be positively related to trait self-control  
109 and self-control resources, which will in turn be related to health behaviors<sup>1</sup>. Figure 1  
110 summarizes our hypotheses.

## 111 2. Method

### 112 Overview

113 Participants were recruited via social media, personal mailing lists, and direct  
114 advertising messages during classes. They were all students at sports and psychology faculties  
115 of three different universities. Data were collected over three weeks (November 2021) by  
116 completion of three cognitive tasks and four questionnaires on Inquisit web version 6.3.2.0  
117 (Computer software) (data hosted by Inquisit, Europe repository) (during a single session  
118 lasting 1h15). All procedures in this study complied with APA ethical principles. Informed  
119 consent was obtained from all participants before the beginning of the study. It should also be  
120 noted that the participants were informed that the online study was anonymous and  
121 confidential: only a self-generated code allowed their identification.

---

<sup>1</sup>No differential hypotheses regarding the role of trait self-control and self-control resources were formulated because of the rare existing evidences about relations between self-control resources and health behaviors. Moreover, we did not formulate differential hypotheses regarding the role of self-control and behaviors, because (a) despite differences, health behaviors considered share similarities (e.g., long-term benefits), as well as unhealthy behaviors share some (e.g., appetitive) (McEachan et al., 2010); and (b) literature stressed that self-control behave in comparable manner on different healthy (e.g., positive relation) and unhealthy behaviors (e.g., negative relation) (de Ridder et al., 2012).

## 122 **Participants and sample size**

123         Three procedures were used for estimating our sample size, and its relevance. The first  
124 procedure aims to estimate a minimal sample size to reach. The second one aims to determine  
125 a stopping rule to fix when data collection will end. The last one aims to compare the sample  
126 size reached and the smallest detectable effect size with a certain power to what the literature  
127 identified in past studies with common features, to consider whether another phase of data  
128 collection is necessary or not. Precisely, first, we used a method specific to structural equation  
129 modeling (MacCallum et al., 1996, 2006a; MacCallum & Hong, 1997) to estimate an a priori  
130 minimum sample size to obtain a fit index, namely RMSEA, within a given range [0.00; 0.08]  
131 (as recommended on the literature, Brown, 2015), with 90% power and  $\alpha = .05$ . Based on  
132 simulations, the minimum sample size was  $N = 26$  (data and code for this estimation are  
133 available at [https://osf.io/hpsjw/?view\\_only=6e28c8307294494e9eec45d2670efd8d](https://osf.io/hpsjw/?view_only=6e28c8307294494e9eec45d2670efd8d))  
134 (MacCallum et al., 2006b). Second, we estimated an a priori maximum sample size by using  
135 the stopping rule based on resource constraints. Because we endorsed no priors regarding an  
136 expected effect size, we decided to recruit as many participants as possible during the running  
137 of the online study. During these three weeks, 535 people logged on. Exclusion criteria were  
138 individuals who: (1) did not consent to participate; (2) completed the study multiple times; (3)  
139 responded incorrectly to seriousness checks (Aust et al., 2013); (4) did not consider regular  
140 physical activity and healthy diet as important for them (i.e., below 2 on a 1-7 goal-  
141 importance scale, Fishbach et al., 2003). In addition, we used the *performance* package  
142 (version 0.10.0) (Lüdecke et al., 2021) for R-Studio (R Core Team, 2021) to identify  
143 observations that were influential on the nine variables of interest (see Measures section).  
144 Specifically, based on a composite score obtained via the application of multiple outlier  
145 detection algorithms (Lüdecke et al., 2021), we excluded participants classified as influential  
146 by at least half of the methods used by this package (the data and code of this data cleaning

147 are available at [https://osf.io/hpsjw/?view\\_only=6e28c8307294494e9eec45d2670efd8d](https://osf.io/hpsjw/?view_only=6e28c8307294494e9eec45d2670efd8d)). The  
148 application of these exclusion criteria in the described order led to a final sample size of  $N =$   
149 385 participants (154 women;  $M_{age} = 19.42$ ,  $SD_{age} = 2.71$ ).

150 Finally, we conducted a sensitivity power analysis with G\*Power 3.1.9.4 (Faul et al.,  
151 2007) to estimate the minimal effect size on the most constrained multiple regression of the  
152 structural equation models. With our convenience sample of  $N = 385$ ,  $n_{predictors} = 5$  (see  
153 Measures), power = .90,  $\alpha = .05$ , the smallest detectable effect size was  $f^2 = .04$ . This is one of  
154 the smallest small-to-medium effect sizes (range = .02 to .15), in line with findings on the  
155 relationships between self-control and health behaviors (Pfeffer & Strobach, 2018).

## 156 **Measures**

### 157 *Independent Variables*

#### 158 **Inhibition**

159 A recent consensual stop-signal task<sup>2</sup> (Verbruggen et al., 2019) was used to assess  
160 inhibition. The task included 216 trials in total, splitted in 3 blocks of 72 trials. A typical trial  
161 started with a central fixation circle presented for 250 ms, followed by display of the stimulus  
162 (a right- or left-pointing white arrow within a circle) until the participant's response. The  
163 instruction was to respond systematically according to the direction indicated by the arrow by  
164 pressing a predefined keyboard button. However, participants had to stop their response (i.e.,  
165 not press the key) if a signal beep was made after the presentation of the arrow. The delay  
166 between the arrow's presentation and the beep was adjusted up or down by 50 ms as a  
167 function of the participant's performance, starting with an initial delay of 250 ms. The delay

---

<sup>2</sup>We preferred the stop-signal task for inhibition because it has been proposed as more closely related to self-control, as it captures more specifically behavioral control, compare to other task such as the Stroop task (supposedly more associated with cognitive inhibition) (Diamond, 2013). We preferred the letter-memory task for updating, because it was the most strongly associated with "updating" latent factor on Miyake et al., (2000) and Miyake & Friedman (2012). We preferred the number-letter task for updating, because it has been proposed as one of the more reliable, and as more appropriate for adults than other task (Diamond, 2013).

168 could be increased up to 1150 ms if the previous signal-stop was successful, and decreased  
169 down to 50 ms if the previous signal stop failed. This delay is referred to as the Stop-Signal  
170 Reaction Time (SSRT) and gives an estimation for response-inhibition latency in  
171 milliseconds. We calculated the SSRT by using the integration method (Verbruggen et al.,  
172 2019). The lower the SSRT, the more difficult it is to stop the go-process, and the higher the  
173 SSRT, the easier it is to stop the go-process. Accordingly, a lower (higher) SSRT integration  
174 means the participant has stronger (weaker) inhibition.

### 175 **Updating**

176 We used a letter memory task to measure updating<sup>2</sup> (Friedman et al., 2008). A series  
177 of letters appeared consecutively in the center of the screen for a duration of 2.5 s for each  
178 letter. Written instructions asked participants to recall, in forward order, the last three letters  
179 after the last letter's disappearance, by selecting the correct letters from a letter matrix  
180 provided. They had to click "blank" if they skipped a particular letter. The number of letters  
181 per series varied randomly through time (5, 7, or 9 letters). In total, 12 measurement trials  
182 were completed (four of each length). Answers were scored as correct even if the three letters  
183 were not recalled in the correct order (Miyake et al., 2000). The more participants were able  
184 to recall letters per trial, the better their updating was considered.

### 185 **Switching**

186 Switching was assessed with a number-letter task<sup>2</sup> (Miyake et al., 2000). This task  
187 involves two categorization tasks, in which character pairs including a letter and a number  
188 (e.g., 3T, 4A) were presented. The participants were asked to categorize the pair depending on  
189 whether the letter was a consonant or vowel (i.e., letter task), or depending on whether the  
190 number was odd or even (i.e., number task). The tasks alternated between categorizing rules  
191 in a clockwise fashion, and thus used predictable location cues in a 2x2 matrix (i.e., the top of  
192 the matrix for letter categorization and bottom of the matrix for number categorization). Odd-

193 numbered trials were set as “switch task” trials and even-numbered trials as “non-switch task”  
194 trials. Participants responded by button press, and the next stimulus was presented 150 ms  
195 after the response. The whole task was composed of 128 trials. The reaction time switch cost  
196 was calculated by assessing the difference between the correct latency of switch trials and  
197 non-switch trials (Miyake et al., 2000). A positive reaction time switch cost indicates a slower  
198 response in switch trials, than in non-switch trials, and conversely. For example, a highly  
199 positive reaction time switch cost indicates low switching. Trials with reaction times under  
200 150 ms and above 2000 ms were excluded from analyses (Rossell & Nobre, 2004;  
201 Schoonbaert et al., 2011).

## 202 *Mediating Variables*

### 203 **Trait Self-Control**

204 Trait self-control was assessed with the 13-item version of the Brief Self-Control Scale  
205 (Tangney et al., 2004). Participants responded to the following instruction: “For each  
206 sentence, choose what suits you best”, on a 7-point Likert scale from 1 (*Don't agree at all*) to  
207 7 (*Completely agree*), with regard to the different items ( $\alpha = .80$ ,  $\omega = .83$ ).

### 208 **Self-Control Resources**

209 Self-control resources were assessed by the subjective vitality scale, as in previous  
210 studies (Forestier et al., 2018). Participants were asked to answer the 5-item questionnaire  
211 (e.g., “At the moment, I feel alive and full of vitality”), with the following instruction: “For  
212 each item, please indicate the general feeling you have experienced over the past 7 days, by  
213 selecting the most appropriate number” on a 7-point Likert scale from 1 (*Don't agree at all*)  
214 to 7 (*Completely agree*) ( $\alpha = .89$ ,  $\omega = .92$ ).

## 215 *Dependent Variables*

### 216 **Physical Activity and Sedentary Activity**

217 Physical activity and sedentary activity were measured using the International Physical  
218 Activity Questionnaire (Craig et al., 2003). Participants were asked to indicate how much  
219 time they had spent doing moderate-intensity and vigorous-intensity physical activities (i.e.,  
220 MVPA), how much time they had spent walking, and how much time they had spent sitting  
221 and/or lying down (i.e., sedentary activity) in minutes, in their daily life over the last 7 days.

### 222 **Healthy and Unhealthy Diet**

223 Healthy and unhealthy diets were assessed using the Healthy Eating Behavior Scale  
224 (Pelletier et al., 2004), composed of two subscales: four items related to a healthy diet (e.g., “I  
225 eat fruit and vegetables”) and the remaining items related to an unhealthy diet (e.g., “I use  
226 white sugar”). Participants indicated their consumption frequency on a 7-item scale ranging  
227 from 1 (*once or twice per month*) to 7 (*more than three times per day*).

## 228 **3. Results**

229 Descriptive statistics and Pearson’s correlation coefficients are provided in the  
230 Supplementary Materials (Tables S1 and S2). Correlations’ table revealed that self-control  
231 resources are significantly correlated with trait self-control ( $r = .21$ ), MVPA ( $r = .27$ ),  
232 sedentary time ( $r = -.14$ ), healthy diet ( $r = .15$ ), unhealthy diet ( $r = -.12$ ) and switching ( $r = -$   
233  $.13$ ). It also stresses that trait self-control is only significantly correlated with healthy and  
234 unhealthy diet ( $r = .21$  and  $r = -.27$ , respectively). The only behaviors significantly correlated  
235 are MVPA and healthy diet ( $r = .25$ ). Among executive functions scores, switching is  
236 significantly correlated with MVPA ( $r = -.10$ ), and inhibition with healthy diet ( $r = -.10$ ).

## 237 **Structural Equation Modelling**

### 238 *Analytical Strategy*

239           The hypotheses were tested using structural equation modeling with the *Lavaan* package  
240 (version 0.6-8, Rosseel, 2012) in R-Studio (RStudio Team, 2015) (the R script, raw data, and  
241 analysis dataset can be found in the Open Science Framework, at  
242 [https://osf.io/hpsjw/?view\\_only=6e28c8307294494e9eec45d2670efd8d](https://osf.io/hpsjw/?view_only=6e28c8307294494e9eec45d2670efd8d)). We used a two-step  
243 approach (Anderson & Gerbing, 1988). The first step is a confirmatory factor analysis (CFA).  
244 Hence, we verified the construct validity of the measurement model to estimate a reliable one,  
245 by examining factor loadings, modification indices and model fit indices. Second, after a  
246 satisfactory fit was achieved for the measurement model, we tested the structural model (i.e.,  
247 the hypothesized relationships between the variables). The results section presents only the  
248 structural models (see Measurement Models in Supplementary Materials). Model fit was  
249 assessed by examining the comparative fit index (CFI), Tucker–Lewis Index (TLI), and root-  
250 mean-square error of approximation (RMSEA), with a satisfactory model having a CFI and a  
251 TLI over 0.90, a RMSEA below 0.05 (Brown, 2015). Finally, after an estimation of the full  
252 hypothetical model, non-significant paths were removed to estimate model’s parsimony  
253 reliability (MacCallum & Austin, 2000). The statistical significance was set at  $\alpha = .05$ . Data  
254 were standardized prior to model estimation. Indirect effects (i.e., mediation) were estimated if  
255 independent variables and mediators were significantly associated with the dependent  
256 variables, with the *RMediation* package (version 1.2.0) (Tofighi & MacKinnon, 2011).

### 257 *Physical Activity and Sedentary Activity Structural Model*

258           The physical activity and sedentary activity structural model contained two latent  
259 variables (trait self-control and self-control resources) and five observed variables (inhibition  
260 score, updating score, switching score, MVPA score, and sedentary activity). Moreover,  
261 compared with the measurement model, we added covariances between the three executive



262 function scores. The full structural equation model yielded good model-fit indices ( $\chi^2$  (130) =  
263 153.43, RMSEA = 0.02, 90% CI [0.00, 0.03], CFI = 0.99, TLI = 0.99). Partially confirming  
264 our hypothesis (H1), only one of the three executive function scores considered was  
265 significantly associated with self-control. Precisely, switching was negatively related to self-  
266 control resources ( $\beta = -.10$ , 95% CI [-.18, -.01],  $p = .02$ ,  $R^2 = .02$ ), evidencing that the higher  
267 the cost, the lower the self-control resources. Contrary to our hypotheses (H2, and H3),  
268 executive function scores were not significantly associated with physical activity and  
269 sedentary activity (i.e., no direct effect). In partial accordance with H4 and H5, only self-  
270 control resources positively predicted MVPA ( $\beta = .31$ , 95% CI [.18, .43],  $p < .001$ ,  $R^2 = .08$ )  
271 and negatively predicted sedentary activity ( $\beta = -.17$ , 95% CI [-.30, -.05],  $p = .01$ ,  $R^2 = .03$ ).  
272 Trait self-control was not significantly associated with healthy or unhealthy behaviors.  
273 Mediation analysis revealed a significant indirect effect of switching on physical activity  
274 (*indirect* = -.03, 95% CI [-.06, -.00],  $p = .02$ ) and sedentary activity (*indirect* = .02, 95% CI  
275 [.00, .04],  $p = .03$ ), suggesting total mediations through self-control resources, which partially  
276 confirm our hypothesis (H6). Figure 2 shows this structural model<sup>3</sup>.

### 277 ***Healthy and Unhealthy Diet Structural Model***

278 The healthy and unhealthy diet structural model contained four latent variables (trait  
279 self-control and self-control resources, and healthy and unhealthy diet scores), three observed  
280 variables (i.e., inhibition, updating, and switching scores), and the covariances between the  
281 three executive function scores. The full structural equation model yielded good model-fit  
282 indices ( $\chi^2$  (188) = 218.35, RMSEA = 0.02, 90% CI [0.00, 0.03], CFI = 0.99, TLI = 0.99).  
283 Switching was still associated with self-control resources, as in the physical and sedentary  
284 activity model. Contrary to H2, and H3, there was no significant direct association between

---

<sup>3</sup>For exploratory purpose, we conducted the model separately on vigorous and moderate physical activity, and on walking. Tables of these models are in Supplementary Materials (S5, S6, S7)

285 executive functions with healthy and unhealthy diets. Mostly according with H4 and H5,  
286 results revealed that trait self-control positively predicted healthy diet ( $\beta = .28$ , 95% CI [.07,  
287 .50],  $p = .01$ ,  $R^2 = .10$ ) and negatively predicted unhealthy diet ( $\beta = -.64$ , 95% CI [-.91, -.36],  
288  $p < .001$ ,  $R^2 = .19$ ), and that self-control resources also positively predicted healthy diet ( $\beta =$   
289 .13, 95% CI [.02, .23],  $p = .02$ ,  $R^2 = .10$ ). Mediation analysis revealed a significant indirect  
290 effect of switching on healthy diet (*indirect* = -.01, 95% CI [-.03, -.00],  $p = .02$ ), suggesting a  
291 total mediation through self-control resources. This partially confirms H6. Figure 3 shows this  
292 structural model. Table 3 summarizes regression coefficients of the two structural models (the  
293 complete tables are available in the Supplementary Materials, Tables S3 and S4).

#### 294 4. Discussion

295 The present study tested the role of executive functions as predictors of individual  
296 differences in trait self-control and self-control resources, which are likely also associated  
297 with health behaviors (physical activity, healthy diet, sedentary activity, and unhealthy diet).  
298 Partially contrary to our hypotheses, we found small evidence supporting the role of executive  
299 functions as predictors of individual differences in trait self-control or self-control resources.  
300 Precisely, switching was only associated with self-control resources. Similarly, no direct role  
301 of these functions regarding healthy and unhealthy behaviors was observed. Nevertheless, in  
302 support of our hypotheses, there was an association between individual differences in self-  
303 control and health behaviors. Individuals with higher self-control resources practiced more  
304 physical activity, ate healthier, and spent less time being sedentary than individuals with  
305 lower self-control resources. It also indicated that individuals with higher trait self-control  
306 adopted a healthier diet, and a less unhealthy one, than individuals with lower trait self-  
307 control. These findings stress the central role of self-control, and especially of self-control  
308 resources, for adopting health behaviors. Finally, we observed three indirect effects of  
309 switching on behaviors, totally mediated by self-control resources. Individuals with higher

310 switching were more physically active, less sedentary, and ate healthier, through higher self-  
311 control resources. These findings pave the way to a more refined understanding of the  
312 predictors of individual differences in self-control resources, and to the ways by which  
313 executive functioning could influence health behaviors.

#### 314 *Relations Between Executive Functions, Self-Control, and Health Behaviors*

315 Partially according with Necka et al. (2018) and Saunders et al. (2018), our study did not find  
316 strong evidence of significant associations between executive functions and self-control.  
317 However, our intention was to examine the contribution of each executive function to self-  
318 control and health behaviors. Indeed, it was proposed that updating and inhibition might be  
319 the most important executive functions for self-control, enabling a better representation of the  
320 goal and a better inhibition for fighting a threatening temptation (Hofmann et al., 2012).  
321 However, our study does not support these suggestions.

322         Suggestions were divided concerning the role of switching. On the one hand, , when  
323 facing a motivational conflict, individuals with high switching could efficiently switch from  
324 an ineffective conflict resolution strategy (e.g., resisting temptation) to a situationally more  
325 effective (e.g., avoiding temptation) (Hofmann et al., 2012). On the other hand, individuals  
326 with high switching could present lower self-control as this switching could promote quick  
327 and efficient disengagement from a goal-oriented mindset to a mindset oriented toward the  
328 pursuit of tempting alternatives (Hofmann et al., 2012). Our results support partially the first  
329 statement. Individuals with high switching are more likely to be able to shift their mind  
330 toward subjective feelings permitting goal-attainment strategies, such as high self-control  
331 resources, useful for efficient self-control act and adaptative motivational-conflict resolution.

332         Our findings also support the absence of evidence in favor of direct relations between  
333 executive functions, trait self-control and physical activity, as observed in some previous  
334 studies. Pfeffer and Strobach (2017) showed that most composite executive functions scores

335 they calculated were not significantly correlated with trait self-control, except for two  
336 switching scores that were, modestly (i.e., task-cueing  $R^2 = .04$ , alternating-runs  $R^2 = .03$ ;  
337 Pfeffer & Strobach, 2017). They also revealed that most executive function scores were not  
338 significant direct predictors of intention-behavior gap, except for updating score. These  
339 findings highlighted no direct relations either between executive functions and trait self-  
340 control or between executive functions and a physical activity, which is consistent with the  
341 current results. Interestingly, Pfeffer et Strobach (2017) found that half of the executive-  
342 function scores they considered (one inhibition, one updating, and one switching score)  
343 moderated the effects of trait self-control on the physical activity intention-behavior gap.  
344 Together with our results, this suggests that updating and inhibition are not direct predictors  
345 of individual differences in self-control and physical activity, but could rather moderate the  
346 relation between self-control and this behavior. However, because the other half of the  
347 executive-function scores they considered (one inhibition score, one updating score, and one  
348 switching score) showed no interaction with self-control to predict physical activity, further  
349 investigations are required.

350 Similarly, the fact that we found no significant direct relations between executive  
351 functions and unhealthy diet were consistent with Hofmann et al. (2009). Indeed, there study  
352 stressed no direct correlation between candy consumption and different components of  
353 inhibition, but these latest (i.e., executive attention, behavioral inhibition, affect regulation)  
354 consistently moderated the relations between automatic affective reactions and candy  
355 consumption. All other things being equal, our study and the aforementioned results  
356 (Hofmann et al., 2009; Necka et al., 2018; Pfeffer & Strobach, 2017) taken together suggest  
357 that inhibition and updating are not direct predictors of self-control or healthy or unhealthy  
358 behaviors, but could moderate relations between affective reactions (e.g., automatic affective  
359 reactions, conscious experience of temptations) and behaviors, and between self-control and

360 health behaviors. Investigations of inhibition and updating as moderators of the relation  
361 between self-control and health behaviors would be of particular interest.

### 362 *Relations Between Self-Control and Health Behaviors*

363 All the relations found between self-control and health behaviors are consistent with  
364 the literature (de Ridder et al., 2011, 2012; Forestier et al., 2018). Specifically, self-control  
365 resources were related to physical activity, sedentary activity, and healthy diet, while trait  
366 self-control was significantly related only to healthy and unhealthy diet. Furthermore, self-  
367 control aspects have different effects depending on the behavior examined, with quite a  
368 similar effect size previously estimated (Forestier et al., 2018). Precisely, self-control  
369 resources were related to health behaviors with a descriptively comparable effect size (i.e.,  $R^2$   
370 = .08 for physical activity, and  $R^2 = .10$  for healthy diet) and a smaller effect size for the  
371 unhealthy behavior ( $R^2 = .03$ ). It suggests that self-control resources are likely efficient to  
372 promote health behaviors, but less effective in preventing the unhealthy ones. Health  
373 behaviors show differences as well as similarities. For example, physical activity, and a  
374 healthy diet are comparable as they require to be initiated, while sedentary activity and an  
375 unhealthy diet are both things that need to be stopped (McEachan et al., 2010). Nevertheless,  
376 physical activity requires more effort to be initiated than a healthy diet (McEachan et al.,  
377 2010), and sedentary activities appear to be attractive because they preserve energy  
378 expenditure (Cheval & Boisgontier, 2021), while unhealthy food seems attractive because of  
379 the immediate pleasure it provides (Appelhans, 2009; Volkow et al., 2011). Another  
380 interesting perspective could be to investigate the influence of behavioral features on self-  
381 control aspects that could be effective or ineffective in the conflict resolution. This  
382 encouraging finding requires further investigations, for example with other studies' design  
383 (see "Strengths, Limitations, and Future Work Perspectives"), or with other instruments to  
384 capture the behaviors, such as accelerometers for physical and sedentary activities, and daily

385 diet diary or questionnaires such as the new Healthy and Unhealthy Eating Behavior Scale  
386 (Guertin et al., 2020) for diet behavior, to confirm the crucial role of self-control resources for  
387 health behaviors.

### 388 **Strengths, Limitations, and Future Work Perspectives**

389         Several limitations of the present study need to be addressed. First, our study sample  
390 consisted of young and relatively healthy students. Psychological determinants driving  
391 behavior maintenance could differ from those driving behavior change (e.g., habit vs. coping  
392 planning). Hence the current results need to be replicated in individuals engaged in health  
393 behavior change processes. Second, the cross-section of the current study is insufficient to  
394 understand the role of within-person variations in executive functions and self-control in daily  
395 fluctuations of health behaviors. Nevertheless, the within-person variations of inhibition seem  
396 more predictive of snack consumption (i.e., unhealthy diet) than individual differences in  
397 inhibition (Powell et al., 2017). In addition, an important part of the variance of self-control  
398 resources is found at the within-person level (i.e., 60%) despite variance at the between-  
399 person level (Smolders et al., 2013). Thus, longitudinal study designs with daily repeated  
400 measures will be required to properly examine the relationships of within-person executive  
401 functions and self-control resource variations with health behavior fluctuations. Third, our  
402 study considered executive functions and self-control as predictors of the overall level of  
403 health behaviors over a week, without measuring the participant's intention to engage in these  
404 behaviors. Despite including only individuals who considered physical activity and healthy  
405 diet important for them, it remains possible that they did not support a particular intention to  
406 engage in healthy behaviors during the week we considered. For example, some participants  
407 may not have practiced physical activity because they did not intend to, rather than because of  
408 low self-control. Future studies could assess intention-behavior gap instead of health  
409 behaviors' global level over a week. In line with previous studies (Pfeffer et al., 2020; Pfeffer

410 & Strobach, 2017), participants could be asked their intention to engage in a certain quantity  
411 of physical activity, to avoid a certain quantity of sedentary activity, to adopt a healthy diet,  
412 and to avoid unhealthy food before and after all the measurements. Then the discrepancy  
413 between intention endorsed and behaviors actually adopted (i.e., intention-behavior gap)  
414 could be assessed to examine the role of executive functions and self-control in reducing this  
415 gap. Finally, the self-reported nature of the self-control and behaviors measures is probably  
416 another limitation of our study. As significant associations were observed between certain  
417 self-reported measures (e.g., self-control resources and physical activity), we cannot exclude  
418 that a part of these associations was because of a common method bias. However, we also  
419 observed association between reaction-time task and self-reported measures (e.g., updating  
420 and self-control resources), and also evidenced absence of significant association between  
421 other self-reported measures (e.g., trait self-control and behaviors). Though, these specific  
422 associations could rather suggest meaningful associations rather a method artifact. Altogether,  
423 direct and conceptual (e.g., different measures and sample) replications could provide  
424 valuable information, and identify the robustness of current (un)significant relations.

425         This study, nevertheless, has several strengths. We first examined the relative and  
426 distinct role of the three executive functions, namely inhibition, updating, and switching, by  
427 using executive tasks according to recent literature (e.g., Verbruggen et al., 2019). We also  
428 tested our hypotheses with structural equation modeling, which increases the reliability of  
429 scores and relations by (a) explicitly assessing the measurement error; (b) estimating latent  
430 variable scores by scoring observed variables rather than other aggregating methods; and (c)  
431 testing a model where a structure (e.g., covariances) could be imposed and assessed as to fit  
432 of the data (Novikova et al., 2013). The final strength of our study is its good power to detect  
433 small effect size (i.e., power = .90 for  $f^2 = .04$ ).

## 434 **Conclusion**

435 No evidence was found supporting executive functions as direct predictors of the four  
436 health behaviors considered. However, the current results support the role of self-control  
437 resources as a potential way to promote physical activity and reduce sedentary activity, and  
438 trait self-control as a likely determinant to increase adoption of a healthy diet and lower that  
439 of an unhealthy diet. In sum, we hypothesize that *cold* executive functions may not explain  
440 individual differences in self-control or health behaviors, and that the aspects of self-control  
441 (trait or state) that are effective in health behavior adoption depend on the behavioral domain.  
442 This study paves the way to longitudinal studies at the within-person level assessing the  
443 effects of *hot* “affective-related” executive functions on trait and self-control resources and  
444 health behaviors. Those effects can be tested by using affectively-charged executive function  
445 in which the neutral stimuli (e.g., arrows, letter or number) are replaced by affective stimuli  
446 such as pictograms of physical or sedentary activity, or of healthy or unhealthy food (e.g., see  
447 Forestier et al., 2022b, p. 27-28, for a discussion). They can also be tested by using self-  
448 control measures specific to the behaviors considered (e.g., self-control resources for  
449 resolving physical-activity related motivational conflict).

## 450 **5. References**

- 451 Allan, J. L., McMinn, D., & Daly, M. (2016). A bidirectional relationship between executive  
452 function and health behavior: Evidence, implications, and future directions. *Frontiers*  
453 *in Neuroscience*, *10*(AUG), 1–13. <https://doi.org/10.3389/fnins.2016.00386>
- 454 Anderson, J. C., & Gerbing, D. W. (1988). Structural Equation Modeling in Practice: A  
455 Review and Recommended Two-Step Approach. *Psychological Bulletin*, *103*(3), 411–  
456 423. <https://doi.org/10.1037/0033-2909.103.3.411>
- 457 Appelhans, B. M. (2009). Neurobehavioral Inhibition of Reward-driven Feeding: Implications  
458 for Dieting and Obesity. *Obesity*, *17*(4), 640–647.  
459 <https://doi.org/10.1038/oby.2008.638>



- 460 Aust, F., Diedenhofen, B., Ullrich, S., & Musch, J. (2013). Seriousness checks are useful to  
461 improve data validity in online research. *Behavior Research Methods*, *45*(2), 527–535.  
462 <https://doi.org/10.3758/s13428-012-0265-2>
- 463 Brown, T. A. (2015). *Confirmatory factor analysis for applied research*. Guilford  
464 publications.
- 465 Cheval, B., & Boisgontier, M. P. (2021). The Theory of Effort Minimization in Physical  
466 Activity. *Exercise and Sport Sciences Reviews*, *49*(3), 168–178.  
467 <https://doi.org/10.1249/JES.0000000000000252>
- 468 Cheval, B., Orsholits, D., Sieber, S., Courvoisier, D., Cullati, S., & Boisgontier, M. P. (2020).  
469 Relationship Between Decline in Cognitive Resources and Physical Activity. In  
470 *Health Psychology* (Issue February). <https://doi.org/10.1037/hea0000857>
- 471 Cipolotti, L., Spanò, B., Healy, C., Tudor-Sfetea, C., Chan, E., White, M., Biondo, F.,  
472 Duncan, J., Shallice, T., & Bozzali, M. (2016). Inhibition processes are dissociable  
473 and lateralized in human prefrontal cortex. *Neuropsychologia*, *93*(September), 1–12.  
474 <https://doi.org/10.1016/j.neuropsychologia.2016.09.018>
- 475 Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E.,  
476 Pratt, M., Ekelund, U., Yngve, A., Sallis, J. F., & Oja, P. (2003). International physical  
477 activity questionnaire: 12-Country reliability and validity. *Medicine and Science in*  
478 *Sports and Exercise*, *35*(8), 1381–1395.  
479 <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>
- 480 de Ridder, D. T. D., Kroese, F., & Gillebaart, M. (2018). Whatever happened to self-control?  
481 A proposal for integrating notions from trait self-control studies into state self-control  
482 research. *Motivation Science*, *4*(1), 39–49. <https://doi.org/10.1037/mot0000062>
- 483 de Ridder, D. T. D., Lensvelt-Mulders, G., Finkenauer, C., Stok, F. M., & Baumeister, R. F.  
484 (2012). Taking stock of self-control: A meta-analysis of how trait self-control relates  
485 to a wide range of behaviors. *Personality and Social Psychology Review*, *16*(1), 76–  
486 99. <https://doi.org/10.1177/1088868311418749>
- 487 Englert, C., & Rummel, J. (2016). I want to keep on exercising but I don't: The negative  
488 impact of momentary lacks of self-control on exercise adherence. *Psychology of Sport*  
489 *and Exercise*, *26*, 24–31. <https://doi.org/10.1016/j.psychsport.2016.06.001>

- 490
- 491 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\* Power 3: A flexible statistical  
492 power analysis program for the social, behavioral, and biomedical sciences. *Behavior*  
493 *Research Methods*, *39*(2), 175–191.
- 494 Fishbach, A., Friedman, R. S., & Kruglanski, A. W. (2003). Leading Us Not Unto  
495 Temptation: Momentary Allurements Elicit Overriding Goal Activation. *Journal of*  
496 *Personality and Social Psychology*, *84*(2), 296–309. [https://doi.org/10.1037/0022-](https://doi.org/10.1037/0022-3514.84.2.296)  
497 [3514.84.2.296](https://doi.org/10.1037/0022-3514.84.2.296)
- 498 Ford, E. S., Zhao, G., Tsai, J., & Li, C. (2011). Low-Risk lifestyle behaviors and all-cause  
499 mortality: Findings from the national health and nutrition examination survey III  
500 mortality study. *American Journal of Public Health*, *101*(10), 1922–1929.  
501 <https://doi.org/10.2105/AJPH.2011.300167>
- 502 Forestier, C., Chanaleilles, M. de, Deschamps, T., & Chalabaev, A. (2022a). Moving forward  
503 with health behaviour change interventions : Considering the plurality of motivational  
504 forces driving health behaviours and motivational conflicts. *PsyArXiv*.  
505 <https://doi.org/10.31234/osf.io/qvrzw>
- 506 Forestier, C., de Chanaleilles, M., Boisgontier, M. P., & Chalabaev, A. (2022b). From ego  
507 depletion to self-control fatigue: A review of criticisms along with new perspectives  
508 for the investigation and replication of a multicomponent phenomenon. *Motivation*  
509 *Science*, *8*(1), 19–32. <https://doi.org/10.1037/mot0000262>
- 510 Forestier, C., Sarrazin, P., Allenet, B., Gauchet, A., Heuzé, J.-P., & Chalabaev, A. (2018).  
511 “Are you in full possession of your capacity?”. A mechanistic self-control approach at  
512 trait and state levels to predict different health behaviors. *Personality and Individual*  
513 *Differences*, *134*, 214–221. <https://doi.org/10.1016/j.paid.2018.05.044>
- 514 Forouzanfar, M. H., Afshin, A., Alexander, L. T., Biryukov, S., Brauer, M., Cercy, K.,  
515 Charlson, F. J., Cohen, A. J., Dandona, L., Estep, K., Ferrari, A. J., Frostad, J. J.,  
516 Fullman, N., Godwin, W. W., Griswold, M., Hay, S. I., Kyu, H. H., Larson, H. J., Lim,  
517 S. S., ... Zhu, J. (2016). Global, regional, and national comparative risk assessment of  
518 79 behavioural, environmental and occupational, and metabolic risks or clusters of  
519 risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015.  
520 *The Lancet*, *388*(10053), 1659–1724. [https://doi.org/10.1016/S0140-6736\(16\)31679-8](https://doi.org/10.1016/S0140-6736(16)31679-8)

- 521 Friedman, N. P., Miyake, A., Young, S. E., DeFries, J. C., Corley, R. P., & Hewitt, J. K.  
522 (2008). Individual Differences in Executive Functions Are Almost Entirely Genetic in  
523 Origin. *Journal of Experimental Psychology: General*, *137*(2), 201–225.  
524 <https://doi.org/10.1037/0096-3445.137.2.201>
- 525 Friese, M., Hofmann, W., & Wiers, R. W. (2011). On taming horses and strengthening riders:  
526 Recent developments in research on interventions to improve self-control in health  
527 behaviors. *Self and Identity*, *10*(3), 336–351.  
528 <https://doi.org/10.1080/15298868.2010.536417>
- 529 Gillebaart, M. (2018). The ‘Operational’ Definition of Self-Control. *Frontiers in Psychology*,  
530 *9*, 1231. <https://doi.org/10.3389/fpsyg.2018.01231>
- 531 Gillebaart, M., Schneider, I. K., & De Ridder, D. T. D. (2016). Effects of Trait Self-Control  
532 on Response Conflict About Healthy and Unhealthy Food. *Journal of Personality*,  
533 *84*(6), 789–798. <https://doi.org/10.1111/jopy.12219>
- 534 Guertin, C., Pelletier, L., & Pope, P. (2020). The validation of the Healthy and Unhealthy  
535 Eating Behavior Scale (HUEBS): Examining the interplay between stages of change  
536 and motivation and their association with healthy and unhealthy eating behaviors and  
537 physical health. *Appetite*, *144*(April 2019), 104487.  
538 <https://doi.org/10.1016/j.appet.2019.104487>
- 539 Hofmann, W., Friese, M., & Roefs, A. (2009). Three ways to resist temptation: The  
540 independent contributions of executive attention, inhibitory control, and affect  
541 regulation to the impulse control of eating behavior. *Journal of Experimental Social*  
542 *Psychology*, *45*(2), 431–435. <https://doi.org/10.1016/j.jesp.2008.09.013>
- 543 Hofmann, W., Friese, M., Schmeichel, B. J., & Baddeley, A. D. (2011). Working memory and  
544 self-regulation. In N. G. P. New York (Ed.), *Handbook of self-regulation* (2nd ed., pp.  
545 204–225). Guilford Press.
- 546 Hofmann, W., Schmeichel, B. J., & Baddeley, A. D. (2012). Executive functions and self-  
547 regulation. *Trends in Cognitive Sciences*, *16*(3), 174–180.  
548 <https://doi.org/10.1016/j.tics.2012.01.006>
- 549 Kotabe, H. P., & Hofmann, W. (2015). On Integrating the Components of Self-Control.  
550 *Perspectives on Psychological Science*, *10*(5), 618–638.  
551 <https://doi.org/10.1177/1745691615593382>

- 552 Lambourne, K. (2006). The relationship between working memory capacity and physical  
553 activity rates in young adults. *Journal of Sports Science & Medicine*, 5(1), 149–153.
- 554 Lee, I. M., Shiroma, E. J., Lobelo, F., Puska, P., Blair, S. N., Katzmarzyk, P. T., Alkandari, J.  
555 R., Andersen, L. B., Bauman, A. E., Brownson, R. C., Bull, F. C., Craig, C. L.,  
556 Ekelund, U., Goenka, S., Guthold, R., Hallal, P. C., Haskell, W. L., Heath, G. W.,  
557 Inoue, S., ... Wells, J. C. (2012). Effect of physical inactivity on major non-  
558 communicable diseases worldwide: An analysis of burden of disease and life  
559 expectancy. *The Lancet*, 380(9838), 219–229. [https://doi.org/10.1016/S0140-  
560 6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)
- 561 Lopez, R. B., Milyavskaya, M., Hofmann, W., & Heatherton, T. F. (2016). Motivational and  
562 neural correlates of self-control of eating: A combined neuroimaging and experience  
563 sampling study in dieting female college students. *Appetite*, 103, 192–199.  
564 <https://doi.org/10.1016/j.appet.2016.03.027>
- 565 Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:  
566 An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal  
567 of Open Source Software*, 6(60), 3139. <https://doi.org/10.21105/joss.03139>
- 568 MacCallum, R. C., & Austin, J. T. (2000). Applications of structural equation modeling in  
569 psychological research. *Annual Review of Psychology*, 51(1), 201–226.  
570 <https://doi.org/10.1146/annurev.psych.51.1.201>
- 571 MacCallum, R. C., Browne, M. W., & Cai, L. (2006). Testing differences between nested  
572 covariance structure models: Power analysis and null hypotheses. *Psychological  
573 Methods*, 11(1), 19.
- 574 MacCallum, R. C., Browne, M. W., & Sugawara, H. M. (1996). Power analysis and  
575 determination of sample size for covariance structure modeling. *Psychological  
576 Methods*, 1(2), 130.
- 577 MacCallum, R. C., & Hong, S. (1997). Power analysis in covariance structure modeling using  
578 GFI and AGFI. *Multivariate Behavioral Research*, 32(2), 193–210.
- 579 McEachan, R. R. C., Lawton, R. J., & Conner, M. (2010). Classifying health-related  
580 behaviours: Exploring similarities and differences amongst behaviours. *British  
581 Journal of Health Psychology*, 15(2), 347–366.  
582 <https://doi.org/10.1348/135910709X466487>

- 583 Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D.  
584 (2000). The Unity and Diversity of Executive Functions and Their Contributions to  
585 Complex ‘Frontal Lobe’ Tasks: A Latent Variable Analysis. *Cognitive Psychology*,  
586 *41*(1), 49–100. <https://doi.org/10.1006/cogp.1999.0734>
- 587 Necka, E., Gruszka, A., Orzechowski, J., Nowak, M., & Wójcik, N. (2018). The  
588 (In)significance of executive functions for the trait of self-control: A psychometric  
589 study. *Frontiers in Psychology*, *9*(JUL), 1–12.  
590 <https://doi.org/10.3389/fpsyg.2018.01139>
- 591 Novikova, S. I., Richman, D. M., Supekar, K., Barnard-Brak, L., & Hall, D. (2013). NDAR:  
592 A model federal system for secondary analysis in developmental disabilities research.  
593 In *International Review of Research in Developmental Disabilities* (Vol. 45, pp. 123–  
594 153). Elsevier. <https://doi.org/10.1016/B978-0-12-407760-7.00003-7>
- 595 Pelletier, L. G., Dion, S. C., Slovinec-D’Angelo, M., & Reid, R. (2004). Why do you regulate  
596 what you eat? Relationship between forms of regulations, eating behaviors, sustained  
597 behavior change, and psychological well-being. *Motivation and Emotion*, *28*(3), 245–  
598 277.
- 599 Pfeffer, I., Englert, C., & Mueller-Alcazar, A. (2020). Perceived stress and trait self-control  
600 interact with the intention–behavior gap in physical activity behavior. *Sport, Exercise,*  
601 *and Performance Psychology*, *9*(2), 244–260. <https://doi.org/10.1037/spy0000189>
- 602 Pfeffer, I., & Strobach, T. (2017). Executive functions, trait self-control, and the intention-  
603 behavior gap in physical activity behavior. *Journal of Sport and Exercise Psychology*,  
604 *39*(4), 277–292. <https://doi.org/10.1123/jsep.2017-0112>
- 605 Pfeffer, I., & Strobach, T. (2018). Behavioural automaticity moderates and mediates the  
606 relationship of trait self-control and physical activity behaviour. *Psychology and*  
607 *Health*, *33*(7), 925–940. <https://doi.org/10.1080/08870446.2018.1436176>
- 608 Powell, D. J. H., McMinn, D., & Allan, J. L. (2017). Does real time variability in inhibitory  
609 control drive snacking behavior? An intensive longitudinal study. *Health Psychology*,  
610 *36*(4), 356–364. <https://doi.org/10.1037/hea0000471>
- 611 R Core Team, R. (2021). *R: A Language and Environment for Statistical Computing*. R  
612 *Foundation for Statistical Computing, 2021.*

- 613 Rabiau, M., Knäuper, B., & Miquelon, P. (2006). The eternal quest for optimal balance  
614 between maximizing pleasure and minimizing harm: The compensatory health beliefs  
615 model. *British Journal of Health Psychology*, *11*(1), 139–153.  
616 <https://doi.org/10.1348/135910705X52237>
- 617 Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling and more.  
618 Version 0.5–12 (BETA). *Journal of Statistical Software*, *48*(2), 1–36.  
619 <https://doi.org/https://www.jstatsoft.org/v48/i02/>
- 620 Rossell, S. L., & Nobre, A. C. (2004). Semantic priming of different affective  
621 categories. *Emotion*, *4*(4), 354.
- 622 Rouse, P. C., Ntoumanis, N., & Duda, J. L. (2013). Effects of motivation and depletion on the  
623 ability to resist the temptation to avoid physical activity. *International Journal of*  
624 *Sport and Exercise Psychology*, *11*(1), 39–56.  
625 <https://doi.org/10.1080/1612197X.2012.717779>
- 626 RStudio Team. (2015). *RStudio: Integrated Development for R*. RStudio, Inc., Boston, MA.  
627 <http://www.rstudio.com/>
- 628 Saunders, B., Milyavskaya, M., Etz, A., Randles, D., & Inzlicht, M. (2018). Reported Self-  
629 control is not Meaningfully Associated with Inhibition-related Executive Function: A  
630 Bayesian analysis. *Collabra: Psychology*, *4*(1), 1–16.  
631 <https://doi.org/10.1525/collabra.134>
- 632 Schoonbaert, S., Holcomb, P. J., Grainger, J., & Hartsuiker, R. J. (2011). Testing asymmetries  
633 in noncognate translation priming: Evidence from RTs and  
634 ERPs. *Psychophysiology*, *48*(1), 74-81.
- 635 Smolders, K. C. H. J., De Kort, Y. A. W., & Van den Berg, S. M. (2013). Daytime light  
636 exposure and feelings of vitality: Results of a field study during regular weekdays.  
637 *Journal of Environmental Psychology*, *36*(1), 270–279.  
638 <https://doi.org/10.1016/j.jenvp.2013.09.004>
- 639 Sniehotta, F. F., Pesseau, J., & Araújo-Soares, V. (2014). Time to retire the theory of planned  
640 behaviour. *Health Psychology Review*, *8*(1), 1–7.  
641 <https://doi.org/10.1080/17437199.2013.869710>

- 642 Tangney, J. P., Baumeister, R. F., & Boone, A. L. (2004). High Self-Control Predicts Good  
643 Adjustment, Less Pathology, Better Grades, and Interpersonal Success. *Journal of*  
644 *Personality*, 72(2), 271–324. <https://doi.org/10.1111/j.0022-3506.2004.00263.x>
- 645 Tofighi, D., & MacKinnon, D. P. (2011). RMediation: An R package for mediation analysis  
646 confidence intervals. *Behavior Research Methods*, 43(3), 692–700.  
647 <https://doi.org/10.3758/s13428-011-0076-x>
- 648 Verbruggen, F., Aron, A. R., Band, G. P. H., Beste, C., Bissett, P. G., Brockett, A. T., Brown,  
649 J. W., Chamberlain, S. R., Chambers, C. D., Colonius, H., Colzato, L. S., Corneil, B.  
650 D., Coxon, J. P., Dupuis, A., Eagle, D. M., Garavan, H., Greenhouse, I., Heathcote,  
651 A., Huster, R. J., ... Boehler, C. N. (2019). A consensus guide to capturing the ability  
652 to inhibit actions and impulsive behaviors in the stop-signal task. *ELife*, 8, 1–26.  
653 <https://doi.org/10.7554/eLife.46323>
- 654 Volkow, N. D., Wang, G.-J., & Baler, R. D. (2011). Reward, dopamine and the control of  
655 food intake: implications for obesity. *Trends in Cognitive Sciences*, 15(1), 37–46.  
656 <https://doi.org/10.1016/j.tics.2010.11.001>

657


**Table 1***Regression Coefficients from Structural Equation Models*

<b>Physical Activity and Sedentary Time Model</b>							
Independent variable	Dependent variable	Estimate	Estimate 95%CI [LL, UL]	Std. Err.	z	p	R <sup>2</sup>
<i>Regression Slopes</i>							
SSRT integration (I)	Self-control resources	0.02	[-0.06, 0.11]	0.04	0.58	.562	.015
Mean correct letters (U)		0.02	[-0.06, 0.10]	0.04	0.47	.638	
RT switch cost (S)		<b>-0.10*</b>	<b>[-0.18, -0.01]</b>	<b>0.04</b>	<b>-2.27</b>	<b>.023</b>	
SSRT integration (I)	Trait self-control	-0.02	[-0.06, 0.03]	0.03	-0.61	.541	.001
Mean correct letters (U)		0.00	[-0.04, 0.05]	0.02	0.18	.860	
RT switch cost (S)		0.00	[-0.04, 0.05]	0.02	0.20	.842	
SSRT integration (I)	MVPA	0.04	[-0.06, 0.14]	0.05	0.81	.418	.081
Mean correct letters (U)		0.00	[-0.09, 0.10]	0.05	0.10	.924	
RT switch cost (S)		-0.07	[-0.17, 0.03]	0.05	-1.44	.150	
Self-control resources		<b>0.31***</b>	<b>[0.18, 0.43]</b>	<b>0.06</b>	<b>4.83</b>	<b>.000</b>	
Trait self-control		0.06	[-0.18, 0.30]	0.12	0.51	.609	
SSRT integration (I)		-0.06	[-0.16, 0.04]	0.05	-1.11	.266	
Mean correct letters (U)	-0.01	[-0.11, 0.09]	0.05	-0.27	.790	.029	
RT switch cost (S)	Sedentary activity	0.04	[-0.06, 0.14]	0.05	0.77		.441
Self-control resources		<b>-0.17**</b>	<b>[-0.30, -0.05]</b>	<b>0.06</b>	<b>-2.71</b>		<b>.007</b>
Trait self-control		-0.02	[-0.27, 0.23]	0.13	-0.16		.874
<i>Fit Indices</i>							
$\chi^2$	153.43						
$\chi^2_{df}$	130.00						
p_ $\chi^2$	.079						
CFI	.99						
TLI	.99						
RMSEA	.02		[0.00, 0.03]				




<b>Healthy and Unhealthy Diet Model</b>						
<i>Regression Slopes</i>						
SSRT integration (I)	Self-control resources	0.03	[-0.06, 0.11]	0.04	0.58	.559
Mean correct letters (U)		0.02	[-0.06, 0.11]	0.04	0.49	.627
<b>RT switch cost (S)</b>		<b>-0.10*</b>	<b>[-0.18, -0.01]</b>	<b>0.04</b>	<b>-2.28</b>	<b>.023</b>
SSRT integration (I)	Trait self-control	-0.02	[-0.07, 0.03]	0.03	-0.61	.542
Mean correct letters (U)		0.00	[-0.05, 0.05]	0.03	0.18	.856
RT switch cost (S)		0.01	[-0.04, 0.06]	0.03	0.24	.813
SSRT integration (I)	Healthy diet	-0.07	[-0.16, 0.01]	0.04	-1.76	.078
Mean correct letters (U)		0.07	[-0.01, 0.16]	0.04	1.73	.084
RT switch cost (S)		0.06	[-0.02, 0.14]	0.04	1.41	.159
<b>Self-control resources</b>		<b>0.13*</b>	<b>[0.02, 0.23]</b>	<b>0.05</b>	<b>2.34</b>	<b>.019</b>
<b>Trait self-control</b>		<b>0.28**</b>	<b>[0.07, 0.50]</b>	<b>0.11</b>	<b>2.62</b>	<b>.009</b>
SSRT integration (I)	Unhealthy diet	0.02	[-0.07, 0.12]	0.05	0.47	.637
Mean correct letters (U)		-0.02	[-0.11, 0.08]	0.05	-0.33	.743
RT switch cost (S)		-0.07	[-0.16, 0.03]	0.05	-1.36	.174
Self-control resources		-0.09	[-0.21, 0.02]	0.06	-1.55	.121
Trait self-control		<b>-0.64***</b>	<b>[-0.91, -0.36]</b>	<b>0.14</b>	<b>-4.56</b>	<b>.000</b>
<i>Fit Indices</i>						
$\chi^2$		218.35				
$\chi^2_{df}$		188.00				
$p_{\chi^2}$		0.06				
CFI		0.99				
TLI		0.98				
RMSEA		0.02	[0.00, 0.03]			

Note. \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ . MVPA = Moderate to vigorous physical activity, SSRT = Stop-Signal Reaction Time, I = Inhibition, U = Updating, S = Switching

**Figure 1***Hypothetical Models*


Note. These are summarized hypothetical models. The full model will test the relations between each executive function, trait and state self-control and each behavior. RT = Reaction Time, SSRT = Stop-Signal Reaction Time, MVPA = Moderate to Vigorous Physical Activity. Variables with a bold line are latent. Variables with normal line are observed. H6 is the hypothesis related to the effects of executive functions on health behaviors mediated by self-control and is not illustrated

**Figure 2***Physical Activity and Sedentary Activity Full Structural Model*

Note. \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ . SSRT = Stop-Signal Reaction Time, RT = Reaction Time, BSCS = Brief Self-Control Scale, SV = Subjective Vitality, MVPA = Moderate to Vigorous Physical Activity. Path darkness level distinguishes significant and non-significant relations.

Figure 3

## Healthy and Unhealthy Diet Full Structural Model



Note. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ . SSRT = Stop-Signal Reaction Time, RT = Reaction Time, BSCS = Brief Self-Control Scale, SV = Subjective Vitality, HBES = Healthy Eating Behavior Scale. Path darkness level distinguishes significant and non-significant relations.

- 1 • Trait self-control and self-control resources are associated with health behaviors
- 2 • Executive functions could have an effect on self-control and health behaviors
- 3 • Self-control resources predicted physical / sedentary activities and healthy diet
- 4 • Trait self-control predicted healthy and unhealthy diet
- 5 • Hot executive functions (affectively charged) seem promising for self-control

Journal Pre-proof

The authors declare no conflict of interests.

Journal Pre-proof