



HAL
open science

Characteristics of lipids body and extracellular matrix in hydrocarbon-rich microalga *Botryococcus braunii* race A and race B strains

Jin J, Stéphanie Dutertre, K. Yoneda, J Legrand, I Suzuki, Watanabe Mm,
Dominique Grizeau, Catherine Dupré

► To cite this version:

Jin J, Stéphanie Dutertre, K. Yoneda, J Legrand, I Suzuki, et al.. Characteristics of lipids body and extracellular matrix in hydrocarbon-rich microalga *Botryococcus braunii* race A and race B strains. Gen2Bio, Mar 2016, Saint-Brieuc, France. hal-04185193

HAL Id: hal-04185193

<https://nantes-universite.hal.science/hal-04185193>

Submitted on 22 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Characteristics of lipids body and extracellular matrix in hydrocarbon-rich microalga *Botryococcus braunii* race A and race B strains

Jin J¹, Dutertre S², Yoneda K³, Legrand J¹, Suzuki I⁴, Watanabe MM⁴, Grizeau D¹, Dupré C^{1*}

¹ LUNAM Université, CNRS, GEPEA, Université de Nantes, UMR 6144, CRTT, 37 bd de l'Université, BP 406, 44602 Saint-Nazaire Cedex, France

² MRic- BIOSIT UMS CNRS 3480 / US INSERM 018 1 - Campus de Villejean, Bât. 8, 2 avenue du Professeur Léon Bernard, 35043 Rennes Cedex, France

³ Graduate School of Life and Environmental Sciences, University of Tsukuba, Ibaraki 305-8572, Japan

⁴ Faculty of Life and Environmental Sciences, University of Tsukuba, Ibaraki 305-8572, Japan

Background


- *Botryococcus braunii* Kützing :
 - ◆ A hydrocarbon producing microalga
 - ◆ Four races (A, B, L and S) according to the chemical structures
 - ◆ Higher hydrocarbons contents in race A and B,
 - ◆ A biomass-derived alternative jet fuel

- ◆ Hydrocarbons are derived from intracellular lipids
- ◆ Hydrocarbons are accumulated extracellularly within a polymerized matrix


Objective

- the cells and colonies morphologies of the two strains (race A and race B) in relation with kinetic parameters of their respective cultures within a chemostat photobioreactor.

Controlled continuous cultures with two strains




- ➔ A-race *B. braunii* strain SAG 30.81
- ➔ B-race *B. braunii* strain Bot 22
- ➔ Chemostat cultures $D_{opt} = 0.23 \text{ day}^{-1}$



Main hydrocarbons according to the strains

SAG 30.81 Alkadiens and Alkatriens

BOT22 Botryococcene $\text{C}_{34}\text{H}_{58}$



Nile Red staining → neutral lipids (yellow)
Chlorophyll → autofluorescence (red)




➔ Chemical analysis confirmed image analysis

Hydrocarbon race	Strain	$\text{g} \cdot \text{m}^{-2} \cdot \text{d}^{-1}$	Content (% dry weight)
Race A	SAG 30,81	0,33	5,1 ±
Race B	BOT 22	2,96	38,3 ± 3,5

No significant hydrocarbon release from cell colonies at low shear stress





Photo Biosit




Race A SAG 30.81

Scale bar : 10µm



Race B Bot 22

Scale bar : 10µm



Conclusions - perspectives

Comparative study of the effect of shear stress on the oils composition and distribution within the extracellular polymeric matrix

- Chromatographic analysis (HPTLC)
- Morphometric analysis of the extracellular liquid hydrocarbons (Confocal microscopy)